

Effects of the littlest Higgs model with T-parity on Higgs boson production at high energy e^+e^- colliders

Chong-Xing Yue and Nan Zhang

Department of Physics, Liaoning Normal University, Dalian 116029, China*

February 2, 2008

Abstract

The Higgs boson production processes $e^+e^- \rightarrow ZH$, $e^+e^- \rightarrow \bar{\nu}_e\nu_e H$, and $e^+e^- \rightarrow t\bar{t}H$ are very important for studying Higgs boson properties and further testing new physics beyond the standard model(*SM*) in the high energy linear e^+e^- collider(*ILC*). We estimate the contributions of the littlest Higgs model with T-parity(*LHT* model) to these processes and find that the *LHT* model can generate significantly corrections to the production cross sections of these processes. We expect the possible signals of the *LHT* model can be detected via these processes in the future *ILC* experiments.

*E-mail: cxyue@lnnu.edu.cn

The discovery and study of Higgs boson is one of the most important goals of present and future high energy collider experiments[1]. The physics of the Higgs boson will be explored by high energy experiments at the upgraded Tevatron collider at Fermilab and in the near future at the large hadron collider at *CERN(LHC)*. The *LHC* will make the first exploration of the TeV energy range, and will be able to discover Higgs boson in the full mass range, provided it exists[2]. After discovery of the Higgs boson at the *LHC*, one of the most pressing tasks is a proper determination of its properties since it is very important to study the mechanism of electroweak symmetry breaking(*EW SB*), the underlying origin of mass, and test the new physics beyond the standard model (*SM*). The *LHC* will be able to finish a few measurement on the couplings of the Higgs boson to fermions and gauge bosons but the most precise measurements will be performed in the clean environment of a future high energy linear e^+e^- collider(*ILC*)[3]. Furthermore, discoveries at the *LHC* may also point to physics scales beyond the reach of the *ILC*, this area could be accessed later by a multi-TeV e^+e^- collider[4].

The main production mechanism of the Higgs boson H at the *ILC* are the Higgs-strahlung process $e^+e^- \rightarrow ZH$ [5] and the WW fusion process $e^+e^- \rightarrow \bar{\nu}_e\nu_e W^*W^* \rightarrow \bar{\nu}_e\nu_e H$ [6]. The cross section for the Higgs-strahlung process decreases as s^{-1} ($\sqrt{s}$ is the center-of-mass energy) and dominates at low energies, while the cross section for the WW fusion process rise as $\log(s/m_H^2)$ and dominates at high energies. The ZZ fusion process $e^+e^- \rightarrow e^+e^- Z^*Z^* \rightarrow e^+e^- H$ can also contribute to the Higgs boson production. However, the cross section is suppressed by an order of magnitude compared to that for the WW fusion process, due to the ratio of the $W^\pm e\nu_e$ coupling to the $Ze\bar{e}$ coupling, $4C_W^2 \simeq 3$. From the production cross sections for the Higgs-strahlung and WW fusion processes, the absolute values of the Higgs couplings to the electroweak gauge bosons Z and W , and also the Higgs couplings to quark and leptons can be determined to a few percent in a model independent way[1,3].

The top quark, with a mass of the order of the electroweak scale, is the heaviest particle yet discovered. The coupling of Higgs boson to top quark pair, which is the largest one among the Yukawa couplings, should play a key role in a theory generating fermion

masses and is particularly sensitive to the underlying physics. Thus, studying the Yukawa coupling $Ht\bar{t}$ is of particular interest. For a light Higgs boson, precise determination of this coupling can be performed at the *ILC* via the associated production process $e^+e^- \rightarrow t\bar{t}H$ [7]. A precision of around 5% can be reached at an *ILC* with $\sqrt{s}=800\sim 1000\text{GeV}$ and the integrated luminosity $\mathcal{L}_{int} \simeq 1000fb^{-1}$ [3,4].

The aim of this letter is to consider the processes $e^+e^- \rightarrow ZH$, $e^+e^- \rightarrow \bar{\nu}_e\nu_e H$, and $e^+e^- \rightarrow t\bar{t}H$ in the context of the littlest Higgs model with T-parity i. e. *LHT* model[8], and see whether the effects of this model on these processes can be detected in the future *ILC* experiments.

To solve the so-called hierarchy or fine-tuning problem of the *SM*, the little Higgs models were proposed as kind of *EW**SB* mechanism accomplished by a naturally light Higgs sector[9]. The key feature of this kind of models is that the Higgs boson is a pseudo-Goldstone boson of a global symmetry which is spontaneously broken at some higher scale f . So far, a number of specific models have been proposed, which differ in the assumed higher symmetry and in the representation of the scalar multiplets. The littlest Higgs(*LH*) model[10] is one of the simplest and phenomenologically viable models, which realizes the little Higgs idea. Most phenomenological analysis about the little Higgs models are given in the context of the *LH* model[11,12].

It has been shown that the *LH* model suffers from severe constraints from precision electroweak measurement, which could only be satisfied by fine-tuning the model parameters[11,13]. To avoid this problem, T-parity is introduced into the *LH* model, which is called *LHT* model[8]. In the *LHT* model, all the *SM* particles are assigned with an even T-parity, while all the new particles are assigned with an odd T-parity, except for the little Higgs partner of the top quark. Thus, the *SM* gauge bosons can not mix with the new gauge bosons, and the electroweak precision observables are not modified at tree level. It has been shown that loop corrections to precision electroweak observables are much small and the scale parameter f can be decreased to 500GeV[8,14]. Thus, the *LHT* model can produce rich phenomenology in the present and future high energy experiments[15,16,17,18].

In the *LHT* model, all the *SM* particles are T-even, while most of the new particles appeared at the TeV scale are T-odd, except for the heavy vector-like top quark T^+ . Thus, the couplings of the electroweak gauge bosons to light fermions are not modified from their corresponding *SM* couplings at tree level. However, since the new T-odd fermions and T-even heavy top quark T^+ are introduced into the *LHT* model, the couplings of the electroweak gauge bosons to top quark and the couplings of the Higgs bosons to ordinary particles are corrected at tree level [15,16]. The expressions of the couplings HVV ($V = W$ or Z), $Ht\bar{t}$, $Zt\bar{t}$, and ZtT^+ , which are related to the processes $e^+e^- \rightarrow \bar{\nu}_e\nu_e H$, $e^+e^- \rightarrow ZH$, and $e^+e^- \rightarrow t\bar{t}H$, can be written as:

$$g_{HV_\mu V_\nu} = \frac{2M_V^2}{\nu} \left[1 - \frac{1}{4} \frac{\nu^2}{f^2} - \frac{1}{32} \frac{\nu^4}{f^4} \right] g_{\mu\nu}; \quad (1)$$

$$g_{Ht\bar{t}} = -\frac{m_t}{\nu} \left[1 - \left(\frac{3}{4} - C_\lambda^2 + C_\lambda^4 \right) \frac{\nu^2}{f^2} \right]; \quad (2)$$

$$g_{Zt\bar{t}}^L = \frac{e}{S_W C_W} \left[\frac{1}{2} - \frac{2}{3} S_W^2 - \frac{C_\lambda^4}{2} \frac{\nu^2}{f^2} \right], \quad g_{Zt\bar{t}}^R = -\frac{e}{S_W C_W} \frac{2}{3} S_W^2; \quad (3)$$

$$g_{Ht\bar{T}^+}^L = -\frac{m_t}{f} S_\lambda^2, \quad g_{Ht\bar{T}^+}^R = \frac{m_t}{\nu} \frac{C_\lambda}{S_\lambda}; \quad (4)$$

$$g_{Zt\bar{T}^+}^L = -\frac{e}{2S_w C_w} C_w^2 \frac{\nu}{f} \sqrt{1 - C_\lambda^4 \frac{\nu^2}{f^2}}, \quad g_{Zt\bar{T}^+}^R = 0. \quad (5)$$

Where $\nu \simeq 246$ is the electroweak scale, $S_W = \sin\theta_W$, θ_W is the Weinberg angle, f is the scale parameter. The mixing parameter $C_\lambda^2 = \frac{\lambda_1^2}{\lambda_1^2 + \lambda_2^2} (S_\lambda^2 = 1 - C_\lambda^2)$, in which λ_1 and λ_2 are the Yukawa coupling parameters.

From above discussions, we can see that the *LHT* model can generate corrections to the production cross sections for the processes $e^+e^- \rightarrow \bar{\nu}_e\nu_e H$ and $e^+e^- \rightarrow ZH$ via the modification of the couplings HWW and HZZ . The value of the relative correction parameters $R_1 = \frac{\sigma^{LHT}(H) - \sigma^{SM}(H)}{\sigma^{SM}(H)}$ and $R_2 = \frac{\sigma^{LHT}(ZH) - \sigma^{SM}(ZH)}{\sigma^{SM}(ZH)}$ are easily calculated, in which $\sigma^{SM}(H)$ and $\sigma^{SM}(ZH)$ represent the corresponding cross sections predicted by the *SM*. From Eq.(1), we can see that there is $R_1 = R_2$, thus we have taken $R = R_1 = R_2$ and plotted R as a function of the scale parameter f in Fig.1.

For the *LH* model, the extra contributions to the process $e^+e^- \rightarrow \bar{\nu}_e\nu_e H$ come from




Figure 1: The relative correction parameter R as a function of the scale parameter f .

the heavy gauge bosons $W_H^\pm$, the modification of the relations among the SM parameters and the precision electroweak input parameters, and the correction terms of the SM $W_e\nu_e$ and HWW couplings[19]. Considering the constraints of the electroweak precision data on the LH model, the scale parameter f should be in the range of 1TeV $\sim$ 3TeV[13]. If we take $f \geq 2\text{TeV}$, the relative correction of the LH model to the production cross section for the process $e^+e^- \rightarrow \bar{\nu}_e\nu_e H$ is smaller than 5%. Thus, the effects of the LHT model on this process are easy to be detected than those of the LH model in the future ILC experiments. The contributions of the LH model to the process $e^+e^- \rightarrow ZH$ mainly come from the heavy photon B_H [20]. For reasonable values of the mixing parameter c' and the mass parameter M_{B_H} , the relative corrections of the LH model to the production cross section for the process $e^+e^- \rightarrow ZH$ can be significantly large. Thus, the contributions of the LHT model to this process can be larger or smaller than those of the LH model.

In the SM , the process $e^+e^- \rightarrow t\bar{t}H$ proceeds mainly through Higgs boson emission off top quarks, while emission from intermediate Z boson plays only a minor role, which has been extensively studied in Ref.[7]. The contributions of the LH model to this

process mainly come from the new gauge bosons B_H and Z_H . In sizable regions of the parameter space preferred by the electroweak precision data, the absolute value of the relative correction parameter $R_3 = \delta\sigma(t\bar{t}H)/\sigma^{SM}(t\bar{t}H)$ can be larger than 5%[21]. For the LHT model, the T-odd new gauge bosons can not contribute to this process at tree level. Thus, the contributions of the LHT model to the production cross section of the process $e^+e^- \rightarrow t\bar{t}H$ only come from the heavy T-even top-quark T^+ and the correction terms to the SM couplings $Zt\bar{t}$, $Ht\bar{t}$, and HZZ .




Figure 2: The relative correction parameter R_3 as a function of the mixing parameter C_λ for four values of the scale parameter f .

Similar to Ref.[21], we can give the production cross section of the process $e^+e^- \rightarrow t\bar{t}H$ in the context of the LHT model. The relative correction parameter R_3 generated by the LHT model is plotted as a function of the mixing parameter C_λ for four values of the scale parameter f in *Fig.2*. One can see from *Fig.2* that, as long as $f \leq 1\text{TeV}$, the absolute value of R_3 is larger than 7%, which might be detected in the future *ILC* experiments.

Since the scale parameter f of the LHT model can be allowed to be lower than 1TeV, this model can produce rich phenomenology in the present and future experiments.

In this letter, we estimate the corrections of the *LHT* model to the production cross sections of the processes $e^+e^- \rightarrow ZH$, $e^+e^- \rightarrow \bar{\nu}_e\nu_e H$, and $e^+e^- \rightarrow t\bar{t}H$, which are very important for studying Higgs properties and test new physics beyond the *SM*. Our numerical results show that, with reasonable values of the free parameters, the *LHT* model can generate significantly contributions to these processes, which might approach the observable threshold of the near-future *ILC* experiments.

Acknowledgments

C. X. Yue would like to thank the **Abdus Salam** International Centre for Theoretical Physics(ICTP) for partial support. This work was supported in part by Program for New Century Excellent Talents in University(NCET-04-0290), the National Natural Science Foundation of China under the Grants No.10475037 and 10675057.

References

- [1] For reviews, See: M. Carena and H. E. Haber, *Prog. Part. Nucl. Phys.* **50** (2003)63; C. Buttar et al. [Standard Model and Higgs Working Group], *hep-ph/0604120*.
- [2] ATLAS Detector and Physics Performance Technical Design Report, <http://atlas.web.cern.ch/Atlas/GROUPS/PHYSICS/TDR/access.html>;
CMS Physics Technical Design Report, <http://cmsdoc.cern.ch/cms/cpt/tdr/>; F. Gianotti and M. L. Mangano, *hep-ph/0504221*.
- [3] T. Abe et al. [American Linear Collider Working Group], *hep-ex/0106057*; J. A. Aguilar-Saavedra et al. [*ECFA/DESY LC* Physics Working Group], *hep-ph/0106315*; K. Abe et al. [*ACFA* Linear Collider Working Group], *hep-ph/0109166*; G. Laow et al., *ILC* Technical Review Committee, second report, 2003, SLAC-R-606.
- [4] E. Accomando et al. [CLIC Physics Working Group], *hep-ph/0412251*.

- [5] J. R. Ellis, M. K. Gaillard, D. V. Nanopoulos, *Nucl. Phys. B***106**(1976)292; B. W. Lee, C. Quigg and H. B. Thacker, *Phys. Rev. D***16**(1977)1519; B. L. Ioffe, V. A. Khoze, *Sov. J. Nucl. Phys.* **9**(1978)50; V. D. Barger et al. *Phys. Rev. D***49**(1994)79;
- [6] D. R. T. Jones and S. T. Petcov, *Phys. Lett. B***84**(1979)440; R. N. Cahn and S. Dawson, *Phys. Lett. B***136**(1984)196; G. L. Kane, W. W. Repko and W. B. Rolnick, *Phys. Lett. B***148**(1984)367; G. Altarelli, B. Mele and F. Pitolli, *Nucl. Phys. B***287**(1987)205; W. Kilian, M. Kramer and P. M. Zerwas, *Phys. Lett. B***373**(1996)135.
- [7] K. J. F. Gaemers and G. J. Gounaris, *Phys. Lett. B***77**(1978)379; A. Djouadi, J. Kalinowski and P. M. Zerwas, *Z. Phys. C***54**(1992)255; S. Dittmaier et al., *Phys. Lett. B***441**(1998)383; S. Dawson and L. Reina, *Phys. Rev. D***57**(1998)5851; *Phys. Rev. D* **59**(1999)054012; *Phys. Rev. D***60**(1999)015003; H. Baer, S. Dawson, L. Reina, *Phys. Rev. D***61**(2000)013002; S. Dittmaier et al., *Phys. Lett. B***478**(2000)247; S. H. Zhu, *hep-ph/0212273*; B. A. Kniehl, F. Madricardo and M. Steinhauser, *Phys. Rev. D***66**(2002)054016; Y. You et al., *Phys. Lett. B***571**(2003)85; G. Belanger et al., *Phys. Lett. B***571**(2003)163; X. H. Wu, C. S. Li and J. J. Liu, *hep-ph/0308012*; A. Denner et al., *Phys. Lett. B***575**(2003)290; *Nucl. Phys. B***680**(2004)85; P. Hafliger, M. Spira, *Nucl. Phys. B* **719**(2005)35; Sun Hao et al., *Phys. Rev. D***71**(2005)075005; Peng Wu et al., *Phys. Lett. B***618**(2005)209.
- [8] H. C. Cheng and I. Low, *JHEP* **0309**(2003)051; *JHEP* **0408**(2004)061; I. Low, *JHEP* **0410**(2004)067.
- [9] For recent review see: M. Schmaltz and D. Tucker-Smith, *Ann. Rev. Nucl. Parti Sci.* **55**(2005)229; T. Han, H. E. Logan, and L. T. Wang, *JHEP* **0601**(2006)099.
- [10] N. Arkani-Hamed, A. G. Cohen, E. Katz, A. E. Nelson, *JHEP* **0207**(2002)034.
- [11] T. Han, H. E. Logan, B. McElrath and L. T. Wang, *Phys. Rev. D***67**(2003)095004.
- [12] M. Perelstein, *hep-ph/0512128*.

- [13] Mu-Chun Chen and S. Dawson, *Phys. Rev. D***70**(2004)015003; R. Casalbuoni, A. Deandrea, M. Oertel, *JHEP* **0402**(2004)032; Chong-Xing Yue and Wei Wang, *Nucl. Phys. B***683**(2004)48; W. Kilian and J. Reuter, *Phys. Rev. D***70**(2004)015004; J. L. Hewett, F. J. Petriello and T. G. Rizzo, *JHEP* **0310**(2003)062; C. Csáki, J. Hubisz, G. D. Kribs, P. Meade, and J. Terning, *Phys. Rev. D***67**(2003)115002; C. Csáki, J. Hubisz, G. D. Kribs, P. Meade, and J. Terning, *Phys. Rev. D***68**(2003) 035009; T. Gregoire, D. R. Smith and J. G. Wacker, *Phys. Rev. D***69**(2004)115008.
- [14] J. Hubisz, P. Meade, A. Noble, and M. Perelstein, *JHEP* **0601**(2006)135.
- [15] J. Hubisz and P. Meade, *Phys. Rev. D***71**(2005)035016.
- [16] C. R. Chen, K. Tobe and C.-P. Yuan, *Phys. Lett. B***640**(2006)263.
- [17] M. Asano et al, *hep-ph/0602157*; A. Martin, *hep-ph/0602206*; A. Birkedal, A. Noble, M. Perelstein, and A. Spray, *Phys. Rev. D***74**(2006)035002 ; C. S. Chen, Kingman Cheung, and T.- C. Yuan, *hep-ph/0605314*.
- [18] H. C. Cheng, I. Low, and L. T. Wang, *Phys. Rev. D***74**(2006)055001; C. O. Dib, R. Rosenfeld, A. Zerwekh, *JHEP* **0605**(2006)074; J. Hubisz, S. J. Lee, and G. Paz, *JHEP* **0606**(2006)041; M. Blanke et al., *hep-ph/0605214*; A. J. Buras, A. Poschenrieder, S. Uhlig, W. A. Bardeen, *hep-ph/0607189*; A. Freitas and D. Wyler, *hep-ph/0609103*.
- [19] Chong-Xing Yue, Wei Wang, Zheng-Jun Zong, Feng Zhang, *Eur. Phys. J. C***42**(2005)331;
- [20] Chong-Xing Yue, Shun-Zhi Wang, Dong-Qi Yu, *Phys. Rev. D***68**(2003)115004.
- [21] Chong-Xing Yue, Wei Wang, Feng Zhang, *Commun. Theor. Phys.* **45**(2006)511.